


MuGard[®]

Mucoadhesive Oral Wound Rinse

How to Obtain MuGard[®]

The *MuGard* Patient Reimbursement and Support Center (MPRSC)

Phone: 888-514-7828 Fax 1: 877-591-2505 Alternate fax: 216-591-2505



The MPRSC works to manage the prescription refill process. The MPRSC is staffed by nurses who are available to help answer questions from HCPs or patients.

*Some restrictions may apply.

Working together to manage the suffering caused by oral mucositis





AMAG Pharmaceuticals®, Inc. can help patients obtain MuGard® when they need it

It is recommended that patients obtain *MuGard* prior to the start of their cancer treatment. *MuGard* is used 4 to 6 times a day throughout therapy.

The AMAG Pharmaceuticals MPRSC works to deliver *MuGard* directly to patients.

To order *MuGard*, follow these two steps:

1. Prescribing physician completes and signs the Prescription Form in the Patient Starter Kit (prescription order form can also be downloaded from MuGard.com).
2. Completed form is faxed to **877-591-2505** or **216-591-2505**.

We'll handle the rest

Remember, one bottle lasts about a week. When completing the Prescription Form, the prescriber should indicate the initial quantity to be dispensed (**typically six 8-oz bottles**). Additional refills may be required depending on the cancer regimen and length of treatment. For more information, contact the MPRSC toll-free at **888-514-7828**.

Important information for patients

The MPRSC will contact patients directly by phone to verify their information. **It is important that patients speak with the MPRSC representative.** In the event that patients miss the call, **it is essential that they call the MPRSC back as soon as possible at 888-514-7828.** Without this return call, timely dispatch of *MuGard* cannot be ensured.

Indication and Important Safety Information

Indication: MuGard® Mucoadhesive Oral Wound Rinse is indicated for the management of oral mucositis/stomatitis (that may be caused by radiotherapy and/or chemotherapy) and all types of oral wounds (mouth sores and injuries), including aphthous ulcers/canker sores and traumatic ulcers, such as those caused by oral surgery or ill-fitting dentures or braces.

Contraindications: *MuGard* is contraindicated in patients with known hypersensitivity to any of the ingredients in the formulation.

Special Precautions for Use: Avoid eating or drinking for at least one hour after rinsing with *MuGard*. After use, replace the bottle cap and tightly seal the bottle. Do not use this product after the expiration date shown on the carton and product label. Do not use this product in patients with known sensitivity to any of the product's ingredients. Dilution of the product prior to use is not recommended.

